HEALING OURSELVES WHILE HEALING OTHERS:

NURSING DURING THE CORONAVIRUS PANDEMIC A Webinar Series with Tools and Resources for Professional Nurses

Donna Gaffney, DNSc, PMHCNS-BC, FAAN





Anne Hofmeyer, PhD, MPHC, RN, MACN

Milagros Elia, APRN-BC





Peg Pipchick, PhD, APN

Milagros Elia, APRN-BC

Donna is a psychotherapist, author and educator, has long addressed a wide range of life-altering experiences in the lives of children and families—loss, trauma, and stress. She has counseled professionals, young people and schools in the aftermath of individual and national tragedies -9/11, Sandy Hook, and Hurricane Katrina. In addition to academic papers, Donna is the author of The Seasons of Grief, Helping Children Grow Through Loss. She taught at Columbia University and holds master's degrees from Teachers College, Columbia University; Rutgers University, and a doctorate from the University of Pennsylvania. Her postdoctoral work includes the Prudential Fellowship for Children and the News at Columbia Journalism School. Donna consults for the New York Life Foundation and the Resilient Parenting for Bereaved Families Program at Arizona State University.

Anne holds an Adjunct appointment with the SONM and is a Visiting Professor at Anglia Ruskin University, Cambridge, UK. She is a member of the Royal College of Nursing (RCN) UK and Australian College of Nursing. Her current research is on translating the social neuroscience of empathy and compassion in the context of culture, networks and leadership in nursing and healthcare. Anne holds a PhD and a Master's Degree in Primary Health Care (palliative care specialty) from Flinders University, Australia. Following completion of her PhD in 2002, she was recruited to the Faculty of Nursing, University of Alberta, Canada in 2003. In 2004, she completed an Intensive Bioethics Course at the Joseph P. & Rose F. Kennedy Institute of Ethics, Georgetown University, Washington DC. Millie is the founder of M. Elia Wellness, LLC, a service which offers Integrative Health Program Design within local cancer survivorship communities, larger healthcare systems and organizations. She is the proud recipient of the (SIO) Society for Integrative Oncology's 2019 Clinician Stakeholder Award for the impact her services have had, and continue to have, on the cancer survivorship community. She has twenty years of experience as a Nurse Practitioner and received her Master's of Arts from NYU in Advanced Practice Nursing. Additionally, she is a certified Health and

Wellness Coach and Yoga Instructor.

Peg is an Advanced Practice Psychiatric Nurse and licensed Marriage and Family Therapist. She works with children, individuals and families to help them become more aware of themselves and others through talking and experiencing their feelings. As a Disaster Crisis Counselor and therapist, Peg has counseled individuals after 9/11, hurricanes Rita, Floyd and Sandy. Peg has served as Adjunct Faculty and Guest lecturer in several nursing programs and taught family therapy at Drew University. As a facilitator for the Recovery and Monitoring Program (RAMP), Institute for Nursing, Peg helped nurses whose practice was impaired by drugs, alcohol or other issues. She has a Masters of Arts from NYU in Psychiatric Nursing, is a Graduate of Blanton-Peale Graduate Institute and earned a PHD from Union Institute and University. Peg has a private practice in Cranford, NJ and is a Certified Holotropic Breathwork Facilitator.

When Bearing Witness is Too Much to Bear



Goals



Charley Mackesy

- Name Trauma and traumatic Responses in yourself, and environment.
- Recognize secondary trauma.
- Identify when to ask for help.

Ē

Stress, Crisis, Trauma



Jason Leung for Unsplash

- **Stress-** causes physical or emotional "tension"; short term or chronic; effects are alleviated when the stressor is removed.
- **Crisis** A temporary disruption of coping and problem-solving skills, but it does *not* necessarily present as a life-threatening experience. Resolved (as well as the resulting state of emotional turmoil and disequilibrium) when the crisis event passes

• Trauma - More extreme version of stressful

- events, they are *perceived* as
- life-threatening,
- evoke negative emotions (fear, helplessness)
- Physical responses and emotions that can last long after the event is over.
- Memory of the traumatic event lingers on.
- "Experiencing repeated or extreme exposure to aversive details of... traumatic events (e.g., first responders; police officers, health care providers)" (APA, 2013, p. 271).



The cortex gives sights and sounds meaningfrom what we have learned in the past

Sights and sounds go to the thalamus (the hub) first, then to the alert center and the cortex

Smell and Touch go directly to the alert center of the brain (amygdala) Smells and touch bring on stronger memories of a frightening event **Processing Memories**

aminalis

7. Locus

ceruleus

tory stimuli

Hypothale

Visual str

Olfactory

The amygdala is the Alert center- lets the body and the brain know danger is present and triggers fight or flight response

RECOLLECTION OF MEMORIES

- Recall is better if the context of recollection resembles the context of the event
- "Increasing levels of arousal DIRECT attention to the central features of the arousing event at the expense of peripheral features."

This is the classic "weapon focus"

Fawcett et al., 2016



FLASHBULB MEMORIES

- A vivid detailed recollection of circumstances of *the receiver* of information that is surprising and important.
 - These are momentous occasions
 - They can change over time



"Memory for a **directly experienced** shocking event is more stable than a flashbulb memory of hearing about the same event."



WHAT IS TRAUMA?

- A shocking, frightening, or dangerous event that often exceeds the standard parameters of the human condition and affects the individual emotionally.
 - Directly experienced
 - Witnessed
 - Learned that the event happened to a family member or friend
 - Experienced first hand repeatedly or extreme exposure to details.

THE ANATOMY OF FEAR

A-The Short Cut All points bulletin to the brain. . before we are aware of the danger!



B- The High Road Information analyzed by the brain- can shut off fear response (experience and memory)



THE ANATOMY OF FEAR . . .



... AND HOW THE BODY RESPONDS

By putting the brain on alert, the amygdala triggers a series of changes in brain chemicals and hormones that puts the entire body in anxiety mode



STRESS-HORMONE BOOST

Responding to signals from the hypothalamus and pituitary gland, the adrenal glands pump out

high levels of the stress hormone cortisol. Too much cortisol shortcircuits the cells in the hippocampus, making it difficult to organize the memory of a trauma or stressful experience. Memories lose their context and become fragmented



RACING HEARTBEAT

The body's sympathetic nervous system, responsible for heart rate and breathing, shifts into overdrive. The heart beats

faster, blood pressure rises and the lungs hyperventilate. Sweat increases, and even the nerve endings on the skin tingle into action, creating goose bumps

FIGHT, FLIGHT OR



FRIGHT The senses become hyperalert, drinking in every detail of the surroundings and looking for potential new threats.

Adrenaline shoots to the muscles, preparing the body to fight or flee

DIGESTION SHUTDOWN

The brain stops thinking about things that bring pleasure, shifting its focus instead to identifying potential dangers. To

ensure that no energy is wasted on digestion, the body will sometimes respond by emptying the digestive tract through involuntary vomiting, urination or defecation

THE BRAIN RESPONDS TO THREAT

 There is one goal:
 To prepare for survival
 -Freezing – an instant check.
 Identify location of danger and if one can escape.





NEXT, FIGHT OR FLIGHT





When the danger passes or one escapes, blood pressure, respiratory rate and hormone flow **return** to normal levels as the body settles into **homeostasis**, or equilibrium, once more.

TRAUMA DISRUPTS . . .

• PHYSIOLOGY

- Tachycardia, increased respirations, dilated pupils, dry mouth, sweating, shaking, dizziness



EMOTIONAL PROCESSING

 A negative emotional state — fear of dying, fear of losing emotional control, emotional distress, horror, anger and disgust.





COGNITIVE PROCESSING

Memory – fragmented, out of sequence
Inability to attend/focus



-Time is distorted



Bovin, M. J., & Marx, B. P., 2010

SUPPRESSING & AVOIDING TRAUMA THOUGHTS

 Research studies show that suppressing memories of any type result in a rebound effect.



 "Although not thinking about painful thoughts seems to be a reasonable way to cope . . . trying to forget actually makes it worse."

Wang, 2020; Wegner, 1994

COVID-19 SOURCES OF ANXIETY

- Access to PPE
- Exposure to COVID-19
- Taking infection home to family
- Unavailability for rapid testing
- Uncertainty of organizational support (personal/family)
- Access to childcare (increased work hours, schools closures)
- Material support (food, hydration, lodging, transportation)
- Ability to provide competent care of deployed to a new clinical settings
- Lack of access to up-to-date and accurate information and communication.



Photo: UK DID

RESPONSES TO TRAUMATIC STRESS

- Initial symptoms are varied, complex, and unstable. They can include
 - exhaustion,
 - Confusion,
 - sadness,
 - anxiety,
 - agitation,
 - numbness,
 - dissociation,
 - disorientation,
 - depression,
 - physical arousal, and
 - blunted affect.



Photo: FEMA

WHAT WE CAN LEARN FROM SARS (2003)

Significant distress in 30-50% health care providers

- Quarantine
- Fear of contagion
- Perceived stigma
- Treating colleagues with SARS
- Concern for family health
- Social isolation

APPROACHES

- Fostering individual resilience
- Fostering organizational resilience



SECONDARY TRAUMATIC STRESS

COLLEAGUES, PEERS AND FAMILY MEMBERS, NOT ONLY ON THE FRONTLINES . . BUT IN THE TRENCHES, CLASSROOMS, HOMES

- Parallels the DSM-5 symptoms of PTSD, that is:
- Intrusive reexperiencing,
- Avoidance,
- Alterations in arousal and reactivity,
- Alterations in cognitions and mood, and dissociation.
- Indirect exposure to traumatic stressors or stress reactions extend beyond the above symptom categories.
 - moral distress,
 - decreased empathy,
 - diminished professional self-efficacy, and
 - feeling stigmatized



Nik Shuliahin for Unsplash

STAYING BALANCED IN THE FACE OF SECONDARY TRAUMA

- Psychological needs related to the trauma situation:
 - Trust/dependency
 - Safety
 - Power
 - Esteem
 - Intimacy
 - Independence
 - Frame of reference



Loic Leray for Unsplash

RISK FACTORS FOR SECONDARY TRAUMA

- A personal trauma history
- Level of education
- Trauma training
- Supervision
- Balance in clinical population

TAKE A BREATH.....



AND ANOTHER BREATH...





RESILIENCE

- "Capacity to cultivate strengths to positively meet the challenges of life." (Silliman, 1994)
- "Ability to bounce back from adversity" (Stuart, 2004)
- "Capacity to bounce forward from adversity, strengthened and more resourceful" (Walsh,1998)



RESILIENCE

- Associated with the ability to cope under adverse circumstances.
- A pattern of adaptive behaviors.
- Not a personality characteristic.
- Not a static or permanent state,
- A dynamic process, associated with—but not identical to personality features.
- A quality found at all levels—from the individual to the family to the work place to the community.



andrzej_b

SELF-ASSESSMENT

"Symptom" identification

Recognize empathic distress and secondary trauma triggers



Identify and use your resources

Review your personal and professional history to the present day



ACTION GOALS

Learn and practice arousal/anxiety reduction skills

Learn grounding skills and containment strategies

Contract with yourself for life enhancement

Resolve barriers to success

Initiate conflict resolution

Implement a supportive aftercare plan

PSYCHOLOGICAL FIRST AID (PFA)

- Reach out to those who need help and provide comfort care.
- Recognize basic needs and support problem-solving.
- Validate feelings and thoughts.
- Provide accurate and timely information.
- Connect individuals with support systems.
- Provide education about stress responses.
- Reinforce strengths and positive coping strategies.



Everly Jr, G. S., & Lating, J. M. 2017

ORGANIZATIONAL RESILIENCE

Hospitals, Clinics, Public Health Organizations, Schools and Universities



1. Establishing reserves

- Material
- Relational
- 2. Effective leadership
- 3. Training and Education
- 4. Supportive, collaborative, interdisciplinary relationships which can provide the basis for formal and informal support during a crisis.
- 5. Characteristics of magnet hospitals
- 6. Organizational justice
 - supervisors take their employees' viewpoints into account,
 - suppress their own biases and deal with subordinates in a fair and truthful manner (relational justice), and
 - fairness

Zachary Keimig for Unsplash

Request	Principal desire	Concerns	Key components of response
Hear me	Listen to and act on health care professionals' expert perspective and frontline experience and understand and address their concerns to the extent that organizations and leaders are able	Uncertainty whether leaders recognize the most pressing concerns of frontline health care professionals and whether local physician expertise regarding infection control, critical care, emergency medicine, and mental health is being appropriately harnessed to develop organization-specific responses	Create an array of input and feedback channels (listening groups, email suggestion box, town halls, leaders visiting hospital units) and make certain that the voice of health care professionals is part of the decision-making process
Protect me	Reduce the risk of health care professionals acquiring the infection and/or being a portal of transmission to family members	Concern about access to appropriate personal protective equipment, taking home infection to family members, and not having rapid access to testing through occupational health if needed	Provide adequate personal protective equipment, rapid access to occupational health with efficient evaluation and testing if symptoms warrant, information and resources to avoid taking the infection home to family members, and accommodation to health care professionals at high risk because of age or health conditions
Prepare me	Provide the training and support that allows provision of high-quality care to patients	Concern about not being able to provide competent nursing/medical care if deployed to new area (eg, all nurses will have to be intensive care unit nurses) and about rapidly changing information/communication challenges	Provide rapid training to support a basic, critical knowledge base and appropriate backup and access to experts Clear and unambiguous communication must acknowledge that everyone is experiencing novel challenges and decisions, everyone needs to rely on each other in this time, individuals should ask for help when they need it, no one needs to make difficult decisions alone, and we are all in this together
Support me	Provide support that acknowledges human limitations in a time of extreme work hours, uncertainty, and intense exposure to critically ill patients	Need for support for personal and family needs as work hours and demands increase and schools and daycare closures occur	Provide support for physical needs, including access to healthy meals and hydration while working, lodging for individuals on rapid-cycle shifts who do not live in close proximity to the hospital, transportation assistance for sleep-deprived workers, and assistance with other tasks, and provide support for childcare needs Provide support for emotional and psychologic needs for all, including psychologic first aid deployed via webinars and delivered directly to each unit (topics may include dealing with anxiety and insomnia, practicing self-care, support for those with greater distress), and provide individual support for those with greater distress
Care for me	Provide holistic support for the individual and their family should they need to be quarantined	Uncertainty that the organization will support/take care of personal or family needs if the health care professional develops infection	Provide lodging support for individuals living apart from their families, support for tangible needs (eg, food, childcare), check-ins and emotional support, and paid time off if quarantine is necessary

Table. Requests From Health Care Professionals to Their Organization During the Coronavirus Disease 2019 Pandemic

PREVENTION

- Psychological
 - Life Balance
 - Relaxation
 - Humor





Photo: Bentley Waters

PREVENTION



- Professional
 - Balance
 - Boundaries
 - Getting support
 - Plans for coping
 - Ongoing education
 - Evaluate healing
 - Skill development

DON'T BE A SPONGE . . .



- When we completely take on other people's suffering as our own, we are at risk.
 - Personal distress, feeling threatened, and overwhelmed.
 - It may lead to burnout.
 - Try to be receptive to other people's feelings *without* absorbing those feelings as your own.

FINDING SANCTUARY

A physical and emotional place of safety and security.

Respite from outside threats.

Offers comfort and familiarity.

Facilitates healing and growth.



Post Traumatic Growth



Taha Mazandarani for Unsplash

"Posttraumatic Growth does not deny the distress associated with highly challenging experiences at the time, and at certain times after. The evidence supporting posttraumatic growth does, however, demonstrate the unique capacity for many people to learn and grow from extreme adversity."

Tedeschi et al., 2018, *Posttraumatic growth: Theory, research, and applications*. Preface.

THE FIVE SENSES GROUNDING EXERCISE

This exercise contains two parts, breathwork and using your five senses.

Start by breathing slowly and deeply, tapping into your own portable tool, one that you can use anytime and anywhere. Breathing slowly allows you to activate the part of your nervous system to help calm and bring you to a place where you can think more clearly and return to the present moment.

The second part of the exercise focuses on the five senses— to help ground us— what you see, hear, touch with your skin, taste and smell. Begin by concentrating on your environment, all that is around you. Focusing helps you feel more centered in your own body. Pay attention to things like shape, textures and colors. Here is an abbreviated exercise using just a one sense, seeing, combined with breath work, it will give you an introduction to the entire exercise.

Start by getting comfortable in your seat. Now begin by taking a slow deep breath through your nose, feeling your belly rise. Slowly turn your head side to side, looking around the room where you are right now. Perhaps look out a window, notice what is far away from you, notice what's close to you. Now notice the different colors and shapes and see the light or lack of light. What captures your attention? Notice a part of your body that you don't usually pay attention to— the soles of your feet, the backs of your calves. If it's comfortable for you, begin to look up and slowly look down continue with your breath, moving through your body. As you slowly move your head notice how your neck feels when you're bringing you head back to center and pause for a moment. Finally notice how this practice makes you feel. Now consider doing the same for what you are hearing, touching, tasting and smelling for the more complete exercise that joins all of our five senses with breath. This grinding exercise can be powerful medicine during potentially traumatic events or anytime that you choose. This is also an easy exercise that children can learn.

https://www.youtube.com/watch?v=7o-oqjiLAOs

When Bearing Witness is Too Much to Bear

American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.). https://doi.org/10.1176/appi.books.9780890425596

Adriaenssens, J., de Gucht, V., & Maes, S. (2012). The impact of traumatic events on emergency room nurses: Findings from a questionnaire survey. *International Journal of Nursing Studies*, 49(11), 1411–1422. doi:10.1016/j.ijnurstu.2012.07.003

Beck, C. T. (2011). Secondary traumatic stress in nurses: A systematic review. Archives of Psychiatric Nursing, 25, 1–10. http://dx.doi.org/10.1016/ j.apnu.2010.05.005

Bonanno, G.A., 2004, 'Loss, trauma, and human resilience: Have we underestimated the human capacity to thrive after extremely aversive events?', *American Psychologist* 59(1), 20–28. https://doi.org/10.1037/0003-066X.59.1.20

Bovin, M. J., & Marx, B. P. (2010, November 22). The Importance of the Peritraumatic Experience in Defining Traumatic Stress. Psychological Bulletin. Advance online publication. doi: 10.1037/a0021353

Brymer, M., Jacobs, A., Layne, C., Pynoos, R., Ruzek, J., & Steinberg, A. (2006). Psychological first aid (PFA) Field Operation Guide.

Czaja, A. S., Moss, M., & Mealer, M. (2012). Symptoms of posttraumatic stress disorder among pediatric acute care nurses. *Journal of pediatric nursing*, *27*(4), 357-365.

Everly Jr, G. S., & Lating, J. M. (2017). *The Johns Hopkins guide to psychological first aid*. JHU Press.

Fawcett, J. M., Peace, K. A., & Greve, A. (2016). Looking down the barrel of a gun: What do we know about the weapon focus effect?. *Journal of Applied Research in Memory and Cognition*, *5*(3), 257-263.

Human Resources. (n.d.) A Personal Strategy for Engaging and Building Your Resilience. University of California San Francisco Human Resources. https://hr.ucsf.edu/hr.php?A=1055&AT=cm&org=c

Jung, H., Jung, S. Y., Lee, M. H., & Kim, M. S. (2020). Assessing the Presence of Post-Traumatic Stress and Turnover Intention Among Nurses Post–Middle East Respiratory Syndrome Outbreak: The Importance of Supervisor Support. Workplace Health & Safety, 216507991989769. doi:10.1177/2165079919897693

Kim, S. J., & Yeo, J. H. (2020). Factors Affecting Posttraumatic Stress Disorder in South Korean Trauma Nurses. *Journal of Trauma Nursing*, *27*(*1*), *50–57*. doi:10.1097/jtn.00000000000482

Komachi, M.K., Kamibeppu, K., Nishi, D. & Matsuoka, Y., 2012, 'Secondary traumatic stress and associated factors among Japanese nurses working in hospital', *International Journal of Nursing Practice* 18(1), 155–163. https://doi.org/10.1111/ j.1440-172X.2012.02014.x

Lavoie, S., Talbot, L., & Mathieu, L. (2011). Post-traumatic stress disorder symptoms among emergency nurses: Their perspective and a 'tailor-made' solution. Journal of Advanced Nursing, 67(7), 1514–1522. doi:10.1111/j.1365-2648.200.05584.x

LeDoux, J. E. (2000). Emotion circuits in the brain. Annual review of neuroscience, 23(1), 155-184.

Luftman, K., Aydelotte, J., Rix, K., Ali, S., Houck, K., Coopwood, T. B., ... Davis, M. (2017). PTSD in those who care for the injured. Injury, 48(2), 293–296. doi:10.1016/j.injury.2016.11.001

Maunder, R. G., Lancee, W. J., Rourke, S., Hunter, J. J., Goldbloom, D., Balderson, K., ... & Fones, C. S. (2004). Factors associated with the psychological impact of severe acute respiratory syndrome on nurses and other hospital workers in Toronto. *Psychosomatic Medicine*, *66*(6), 938-942.

Maunder, R. G., Leszcz, M., Savage, D., Adam, M. A., Peladeau, N., Romano, D., ... & Schulman, R. B. (2008). Applying the lessons of SARS to pandemic influenza. *Canadian Journal of Public Health*, 99(6), 486-488.

Mealer, M., Jones, J., & Meek, P. (2017). Factors affecting resilience and development of posttraumatic stress disorder in critical care nurses. *American Journal of Critical Care*, 26(3), 184-192.

Ogino, Y., Kawamichi, H., Kakeda, T., & Saito, S. (2019). Exploring the neural correlates in adopting a realistic view: A neural structural and functional connectivity study with female nurses. *Frontiers in human neuroscience*, 13.

Panagioti, M., Gooding, P. A., Taylor, P. J., & Tarrier, N. (2014). Perceived social support buffers the impact of PTSD symptoms on suicidal behavior: Implications into suicide resilience research. Comprehensive Psychiatry, 55, 104–112. doi:10.1016/j. comppsych.2013.06.004

Salmon, G., & Morehead, A. (2019). Posttraumatic Stress Syndrome and Implications for Practice in Critical Care Nurses. *Critical Care Nursing Clinics*, *31*(4), 517-526.

Schacter, D. L. (2008). *Searching for memory: The brain, the mind, and the past*. Basic Books.

Shalev, A.Y. (n.d.). Treating Survivors in the Acute Aftermath of Traumatic Events. National Center for PTSD.

https://www.ptsd.va.gov/professional/treat/txessentials/tx_survivors_trauma.asp#two

Shalev, A.Y. (2002). Treating Survivors in the Immediate Aftermath of Traumatic Events. In Rachel Yehuda (Ed.), *Treating Trauma Survivors with PTSD* (157-188). Washington DC: American Psychiatric Publishing, Inc.

Shanafelt, T., Ripp, J., & Trockel, M. (2020). Understanding and Addressing Sources of Anxiety Among Health Care Professionals During the COVID-19 Pandemic. *JAMA*. doi: 10.1001/jama.2020.5893

Sprang, G., Ford, J. D., Kerig, P. K., & Bride, B. (2018). Defining secondary traumatic stress and developing targeted assessments and interventions: Lessons learned from research and leading experts. Traumatology. Advance online publication. http://dx.doi.org/10.1037/ trm0000180

Tedeschi, R. G., Shakespeare-Finch, J., Taku, K., & Calhoun, L. G. (2018). *Posttraumatic growth: Theory, research, and applications*. Routledge.

Van der Kolk, B. A., McFarlane, A. C., & Weisaeth, L. (2007). Trauma Stress: The effects overwhelming experiences on mind, body and society. *The Guildford Process, New York, NY*.

Walsh, F. (2007). Traumatic loss and major disasters: Strengthening family and community resilience. *Family process*, *46*(2), 207-227.

Wang, D., Hagger, M. S., & Chatzisarantis, N. L. (2020). Ironic Effects of Thought Suppression: A Meta-Analysis. *Perspectives on Psychological Science*, 1745691619898795.

Wegner, D. M., & Gold, D. B. (1995). Fanning old flames: emotional and cognitive effects of suppressing thoughts of a past relationship. *Journal of personality and social psychology*, *68*(5), 782.

Yehuda R. Discrepancy between theory, research and practice. (1999, September) Conference presentation at Mt. Sinai Medical Center; New York, New York.



Psychological First Aid for Health Care Providers

https://omh.ny.gov/omhweb/disaster resources/pfa/healthcare.pdf



Let us know how you are doing!

HealingOurselves@DonnaGaffneyDNSc.com



Photo: Jason Rosewell for Unsplash